



| | | |
|---|--|--|
| FORM PTO-1390 DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE (REV 1-98) | | ATTORNEY'S DOCKET NO. 11362.0034.PCUS00; (INNS.034) |
| TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371 | | U.S. APPLICATION NO. (If known, see 37 CFR 1.5) |
| | | 09/786481 |
| INTERNATIONAL APPLICATION NO. PCT/EP99/06592 | INTERNATIONAL FILING DATE 7 September 1999 | PRIORITY DATE CLAIMED 8 September 1998 |
| TITLE OF INVENTION TAU AS A MARKER FOR EARLY CNS DAMAGE | | |
| APPLICANT(S) FOR DO/EO/US Frank HULSTAERT; Eugene VANMECHELEN; Hugo VANDERSTICHELE; | | |
| Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information: | | |
| <ol style="list-style-type: none"> <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. <input type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1). <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)). <ol style="list-style-type: none"> <input checked="" type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). <input checked="" type="checkbox"/> has been transmitted by the International Bureau. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371(c)(2)). <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)). <ol style="list-style-type: none"> <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). <input type="checkbox"/> have been transmitted by the International Bureau. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. <input checked="" type="checkbox"/> have not been made and will not be made. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). | | |
| Items 11 to 16 below concern document(s) or information included: | | |
| <ol style="list-style-type: none"> <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. <input checked="" type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. <input checked="" type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. <input type="checkbox"/> A substitute specification. <input checked="" type="checkbox"/> A change of power of attorney and/or address letter. <input checked="" type="checkbox"/> Other items or information: Postcard, Fee Calculation Sheet (in duplicate); | | |

| | |
|--|---|
| CERTIFICATE OF EXPRESS MAILING | |
| NUMBER | EL521279540US |
| DATE OF DEPOSIT | MARCH 5, 2001 |
| SIGNATURE (Jennifer L. Purnell) |  |
| This paper or fee is being deposited with the United States Postal Service "EXPRESS MAIL POST OFFICE TO ADDRESSEE" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to: Commissioner for Patents, Washington, DC 20231. | |

| | | | | | |
|---|--------------|---|------------|---|---------|
| U.S. APPLICATION NO. <u>097786481</u> (if known, see 37 CFR 1.51) | | INTERNATIONAL APPLICATION NO. PCT/EP99/06592 | | ATTORNEY'S DOCKET NUMBER 11362.0034.PCUS00; INNS:034 | |
| 17. <input checked="" type="checkbox"/> The following fees are submitted: Basic National Fee (37 CFR 1.492(a)(1)-(5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1,000.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$860.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$760.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$690.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$ 100.00 ENTER APPROPRIATE BASIC FEE AMOUNT = \$ 860.00 | | | | CALCULATIONS PTO USE ONLY | |
| Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)). | | | | \$ 0.00 | |
| Claims | Number Filed | Number Extra | Rate | | |
| Total Claims | 15-20 = | | x \$ 18.00 | \$ | |
| Independent Claims | 4-3 | 1 | x \$ 80.00 | \$ | 80.00 |
| Multiple dependent claim(s) (if applicable) | | | | \$ | 270.00 |
| TOTAL OF ABOVE CALCULATIONS = | | | | \$ | |
| Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity statement must also be filed. (NOTE: 37 CFR 1.9, 1.27, 1.28) | | | | \$ | .00 |
| SUBTOTAL = | | | | \$ | 1210.00 |
| Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)). | | | | \$ | .00 |
| TOTAL NATIONAL FEE = | | | | \$ | 1210.00 |
| Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property) | | | | \$ | .00 |
| TOTAL FEES ENCLOSED = | | | | \$ | |
| | | | | Amount to be refunded: | \$.00 |
| | | | | charged | \$.00 |
| a. <input type="checkbox"/> A check in the amount of \$_____ to cover the above fees is enclosed. Applicants request a Notice of Missing Parts to satisfy the filing of the Declaration and Preliminary Amendment. b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. <u>01-2508/11362.0034.PCUS00</u> in the amount of <u>\$1210.00</u> to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>01-2508/11362.0034.PCUS00</u> . A duplicate copy of this sheet is enclosed. | | | | | |
| NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status. | | | | | |
| SEND ALL CORRESPONDENCE TO: Patricia A. Kammerer HOWREY SIMON ARNOLD & WHITE, LLP 750 Bering Drive Houston, TX 77057 (713) 787-1400 | | | | | |
|  SIGNATURE | | | | | |
| PATRICIA A. KAMMERER NAME | | | | | |
| <u>29,775</u> REGISTRATION NUMBER | | | | | |

09/786481

528 Rec PCT/PTO 05 MAR 2001

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:

FRANK HULSTAERT**EUGEN VANMECHELEN****HUGO VANDERSTICHELE**

Serial No.:

Filed:

For: **TAU AS A MARKER FOR EARLY CNS
DAMAGE**

§

§

§ Group Art Unit: **Unknown**

§

§ Examiner: **Unknown**

§

§ Atty. Dkt. No.: **11362.0034.PCUS00**

§

§ **INNS:034---**

§

§ History: **International Application No.:**

§

§ **PCT/EP99/06592**

§


§ **Filing Date: 7 September 1999****PRELIMINARY AMENDMENT AND PRIORITY NOTICE**

Box PCT

Commissioner for Patents

Washington, D.C. 20231

Sir:

| | |
|--|----------------------|
| CERTIFICATE OF EXPRESS MAIL | |
| NUMBER | EL521279540US |
| DATE OF DEPOSIT | March 6, 2001 |
| I hereby certify that this paper or fee is being deposited with the United States Postal Service "EXPRESS MAIL POST OFFICE TO ADDRESSEE" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to: Commissioner for Patents, Washington, D.C. 20231. | |
|  Jennifer L. Purnell | |

Please amend this application as follows:

IN THE SPECIFICATION:

On page 1, line 2, please add a new paragraph following the Title:

--This application is a §371 national stage filing of PCT/EP99/06592, filed 7 September 1999 (published in English on 16 March 2000 as WO 00/14546) and claiming priority to EP 98870190.0 filed 8 September 1998.--

At page 4, line 1, please add a Summary Section:

--SUMMARY OF THE INVENTION

The present invention relates to a method for the early detection and/or quantification of CNS damage in an individual, said CNS damage being caused by space-occupying lesions of the CNS, by invasion of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents or by a combination of these mechanisms. This method comprises the step of determining and/or quantifying the level of tau in an individual and comparing it to the level of tau in control healthy individuals.--

At page 5, line 27, please add a new Section:

--BRIEF DESCRIPTION OF THE FIGURES

Figure 1. Tau values at diagnosis, before any treatment was given: 1. Control children; 2. AML; 3. AML-CNS+; 4. CML; 5. MDS; 6. B-NHL; 7. Non-B-ALL; 8. Non-B-ALL CNS+; 9. VHR non-B-ALL.

Figure 2. CSF-tau levels in seven patients after acute ischemic stroke. The CSF samples were collected at intake (day 0-1), day 2-3, day 7-8, day 21-22 (3 weeks) and day 90-110 (3 months).

Figure 3. CSF-tau levels at the time of maximal release in relation to the size of the infraction as measured by CT scan.--

At page 20, please delete the section entitled "Figure Legends" on lines 5-15.

At page 26, line 13 the text should refer to table 4 instead of table 5, therefore please replace the paragraph on lines 11 - 16 to read:

--Tau levels in the CSF samples of patients with possible CNS damage caused by space occupying lesions, by invasion or metastasis, by bleeding, infarction or ischemia or by organisms and tau levels in CSF samples of control subjects are show in table 4. The group of patients that had possibly contracted CNS damage by space occupying lesions, by invasion or metastasis, by bleeding, by infarction or ischemia, or by organisms showed an overall higher tau level than the neurological control patients ($p = 0.022$, two sided Mann Whitney test).--

On a new page following the Claims Section, please add the Abstract found on the coversheet:

--ABSTRACT

The present invention provides a new method for the early diagnosis of CNS damage in an individual, said CNS damage being caused by space-occupying lesions of the CNS, by invasion or metastasis of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents, or by a combination of these mechanisms. This new method comprises the step of determining and/or quantifying the level of tau in said individual and comparing it to the level of tau in control healthy individuals.--

IN THE FIGURES

Please replace original Fig. 2 with the attached corrected Fig. 2, where the legend has been changed from "Patent number" to read --Patient number--.

IN THE CLAIMS:

Please cancel claims 12 and 13.

Please amend claims 5 - 11, 15 and 16 as follows:

5. (Amended) A method according to [any of claims 1 to 4] claims 1 or 2 in which the space-occupying lesion of the CNS is a primary brain tumor, benign or malignant, brain metastasis, or a subdural haematoma.
6. (Amended) A method according to [any of claims 1 to 4] claims 1 or 2 in which the invasion or metastasis of the CNS is by leukemia, lymphoma or breast cancer.
7. (Amended) A method according to [any of claims 1 to 4] claims 1 or 2 in which the organisms are bacteria or viruses causing encephalitis or meningitis.
8. (Amended) A method according to [any of claims 1 to 4] claims 1 or 2 in which the anoxia or ischemia is caused by stroke, by cerebral infarction, by cerebral hemorrhage, by thrombosis, by perinatal asphyxia, by Binswanger disease or by vasculitis.
9. (Amended) A method according to [any of claims 1 to 4] claims 1 or 2 in which the chemical agent is gene therapy, pharmaceuticals, chemotherapy or exposure to chemical compounds.
10. (Amended) A method according to [any of claims 1 to 4] claims 1 or 2 in which the physical agent is a trauma, stroke, intracranial pressure or radiation.

11. (Amended) A method according to [any of claims 1 to 10] claims 1 or 2 in which CNS damage is detected and/or quantified in order to evaluate the effect of a certain treatment on said CNS damage.

15. (Amended) A kit according to claim [14 for use in any method according to claims 1 to 11] 16, wherein the marker is tau.

16. (Amended) A kit according to [claims 13 and/or] claim 14 characterised in that said kit comprises:

- a monoclonal antibody (primary antibody) which forms an immunological complex with an epitope of tau;
- a secondary antibody
 - which can be a monoclonal antibody recognising an epitope of the tau-primary antibody complex, but not recognising the primary antibody alone, or
 - which can be a polyclonal antibody recognising an epitope of the tau-primary antibody complex but not recognising the primary antibody alone, with said polyclonal antibody being preferably purified by immunoaffinity chromatography using immobilized tau or immobilized tau-primary antibody complex;
- a marker either for specific tagging or coupling with said secondary antibody,

- appropriate buffer solutions for carrying out the immunological reaction between the primary antibody and the test sample, between the secondary antibody and the tau-primary antibody complex and/or between the secondary antibody and the marker;
- [possibly,] optionally, for standardisation purposes, a purified protein or a synthetic peptide containing one or more tau epitopes.

REMARKS

The specification has been amended to claim priority and add reference to an earlier filed PCT application as required under 37 C.F.R. § 1.78(a)(2). Further, the specification has been amended to reflect the section headings and organization format preferred by the United States Patent and Trademark Office. Support for the Summary of the Invention is found at page 5, line 31 bridging page 6, line 5. Also attached is an Abstract on a new page.

In this amendment, claims 12 and 13 are cancelled and claims 5-11, 15 and 16 are amended to modify the recitation of multiple dependency. The amendment to claim 9 finds support in the Specification at page 7, lines 5 and 6. The amendment to claim 15 finds support in cancelled claim 12. Claims 1-11 and 14-17 are now pending. A clean set of the pending claims is attached hereto.

It is believed that no fee is due. Should any fees under 37 C.F.R. §§ 1.16 to 1.21 be required for any reason relating to the enclosed materials, the Commissioner is authorized to deduct said fees from Deposit Account No. 01-2508/11362.0034.PCUS00.

CONCLUSION

In view of the foregoing amendments, applicants respectfully submit the claims are in proper form and condition for allowance. Applicants request that the claims be allowed and the application advanced to issue.

The Examiner is invited to contact the undersigned attorney at (713) 787-1438 with any questions, comments or suggestions relating to the referenced patent application.

Respectfully submitted,



Patricia A. Karhmerer
Reg. No. 29,775
Attorney for Assignee
INNOGENETICS N.V.

HOWREY SIMON ARNOLD & WHITE, LLP
750 Bering Drive
Houston, Texas 77057-2198
(713) 787-1400

Date: March 5, 2001

PENDING CLAIMS FOLLOWING PRELIMINARY AMENDMENT

1. A method for the early detection and/or quantification of CNS damage in an individual, said CNS damage being caused by space-occupying lesions of the CNS, by invasion or metastasis of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents or by a combination of these mechanisms, said method comprising the step of determining the level of tau in said individual and comparing it to the level of tau in control healthy individuals.
2. A method for the early in vitro detection and/or quantification of CNS damage in an individual, said CNS damage being caused by space-occupying lesions of the CNS, by invasion or metastasis of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents or by a combination of these mechanisms, said method comprising the steps of:
 - obtaining a sample from said individual,
 - determining the level of tau in said sample and comparing it to the level of tau in control healthy individuals.
3. A method according to claim 2 in which the sample is taken from the cerebrospinal fluid of the individual.
4. A method according to claim 2 in which the sample is taken from the blood derivatives of the individual.

5. (Amended) A method according to claims 1 or 2 in which the space-occupying lesion of the CNS is a primary brain tumor, benign or malignant, brain metastasis, or a subdural haematoma.

6. (Amended) A method according to claims 1 or 2 in which the invasion or metastasis of the CNS is by leukemia, lymphoma or breast cancer.

7. (Amended) A method according to claims 1 or 2 in which the organisms are bacteria or viruses causing encephalitis or meningitis.

8. (Amended) A method according to claims 1 or 2 in which the anoxia or ischemia is caused by stroke, by cerebral infarction, by cerebral hemorrhage, by thrombosis, by perinatal asphyxia, by Binswanger disease or by vasculitis.

9. (Amended) A method according to claims 1 or 2 in which the chemical agent is gene therapy, pharmaceuticals, chemotherapy or exposure to chemical compounds.

10. (Amended) A method according to claims 1 or 2 in which the physical agent is a trauma, stroke, intracranial pressure or radiation.

11. (Amended) A method according to claims 1 or 2 in which CNS damage is detected and/or quantified in order to evaluate the effect of a certain treatment on said CNS damage.

14. A kit for the early diagnosis of CNS damage in an individual, said CNS damage being caused by space-occupying lesions of the CNS, by invasion of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents, or by a combination of these mechanisms, comprising a tool for the detection of tau.

15. (Amended) A kit according to claim 16, wherein the marker is tau.

16. (Amended) A kit according to claim 14 characterised in that said kit comprises:

- a monoclonal antibody (primary antibody) which forms an immunological complex with an epitope of tau;
- a secondary antibody
 - which can be a monoclonal antibody recognising an epitope of the tau-primary antibody complex, but not recognising the primary antibody alone, or
 - which can be a polyclonal antibody recognising an epitope of the tau-primary antibody complex but not recognising the primary antibody alone, with said polyclonal antibody being preferably purified by immunoaffinity chromatography using immobilized tau or immobilized tau-primary antibody complex;
- a marker either for specific tagging or coupling with said secondary antibody,
- appropriate buffer solutions for carrying out the immunological reaction between the primary antibody and the test sample, between the secondary

antibody and the tau-primary antibody complex and/or between the secondary antibody and the marker;

- optionally, for standardisation purposes, a purified protein or a synthetic peptide containing one or more tau epitopes.

17. A method to screen or monitor the effect of compounds which prevent or treat CNS damage comprising the step of determining the level of tau and comparing it to the level of tau in a control sample.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218

ABSTRACT

The present invention provides a new method for the early diagnosis of CNS damage in an individual, said CNS damage being caused by space-occupying lesions of the CNS, by invasion or metastasis of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents, or by a combination of these mechanisms. This new method comprises the step of determining and/or quantifying the level of tau in said individual and comparing it to the level of tau in control healthy individuals.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216

09/786481




Figure 2

09/786489




Figure 2

TAU AS A MARKER FOR EARLY CNS DAMAGE

FIELD OF THE INVENTION

5

The present invention relates to the field of CNS damage. The present invention relates to a new method for the early diagnosis of CNS damage by detection and/or quantification of tau.

10 BACKGROUND OF THE INVENTION

Damage of the central nervous system (CNS damage) is caused by various inducing agents among which different disease processes, physical or chemical agents, anoxia and ischemia. Disease processes include space occupying lesions, invasion or metastasis of the brain caused by different malignancies and/or infection by a number of organisms.

15

Tumors of the CNS, may originate locally (primary tumors) or may spread to the CNS (metastases). Primary tumors arise from glial cells (astrocytoma, oligodendroglioma, glioblastoma), ependymal cells (ependymoma) or supporting tissue (meningioma, schwannoma, papilloma of the choroid plexus). In childhood, tumors arise from more primitive cells (medulloblastoma, neuroblastoma, chordoma). Malignant astrocytoma or glioblastoma is the most common type of primary tumor in adults over age 20. Both benign and malignant primary CNS tumors are capable of producing neurologic impairment.

20

Leukemia is the most common type of cancer in children. During the last twenty years, the survival of children with leukemia has improved markedly based on the routine use of intensive chemotherapy alone or as combined treatment (radiotherapy and chemotherapy). Currently, the estimated overall 10-year survival rate is around 75%. Given the increasing number of childhood leukemia survivors, concern has arisen about long-term effects of anti-cancer chemotherapy and/or radiotherapy resulting in possible damage to the central nervous system and the need for an early quantitative determination of this CNS damage is increasing.

25

30 Bacterial meningitis may be defined as an inflammation in response to bacterial infection of the pia-arachnoid and the fluid residing in the space which it encloses and also of the fluid in the

ventricles of the brain. The incidence of bacterial meningitis is between 4.6 and 10 cases per 100000 persons per year. *H. influenzae* is the most frequent cause, followed by *N. meningitidis* and *S. pneumoniae*. Once developed, characteristic features of bacterial meningitis include an increase in intracranial pressure, disruption of the blood-brain barrier, cerebral edema, and alterations in cerebral blood flow. The longer the duration of meningitis and the less effective the treatment, the greater the chances that complications and neurologic residua will develop. Approximately 10 percent of infants and children who have bacterial meningitis will be left with persistent unilateral or bilateral sensory hearing loss. Approximately 30 percent of children who have had bacterial meningitis later will turn out to have subtle learning deficits (Wilson et al., 1991).

Viruses can also affect the central nervous system in a variety of ways resulting in a distinction between viral meningitis, viral encephalitis, myelitis and CNS diseases due to slow virus infection. Other conditions that may cause CNS damage are chemical agents such as pharmaceuticals, chemotherapy or exposure to chemical compounds, and physical agents. Head injuries are frequent in industrialised countries, affecting many patients in the prime of life. To appreciate the medical and social magnitude of this problem it needs only to be recognised that almost 10 million Americans have head injuries yearly, about 20 percent serious enough to cause brain damage. Another cause for CNS damage may be anoxia or ischemia. Anoxic-ischemic encephalopathy is a common and often disastrous condition, caused by a lack of oxygen to the brain, resulting from hypotension or respiratory failure. Acute ischemic stroke causes neuronal damage and is a major cause of neurological handicap in western society. Perinatal asphyxia may be associated with CNS damage as well. To date, clinical, electroencephalographic and neuroradiologic evaluation, together with cerebral blood flow studies are the most readily available methods. However, early and accurate evaluation of the severity of brain damage after a hypoxic-ischemic event, remains one of the most difficult problems in neonatal care.

For the detection of CNS invasion in leukemic children current diagnostic procedures include lumbar puncture, eye fundoscopy and brain imaging (Raichle, 1998). However, these diagnostic methods only allow detection of the CNS damage in a more advanced stage while already ongoing CNS damage in the early stages may be missed by these methods. Therefore, there is a need for additional diagnostic methods that allow early detection of CNS damage.

A number of neurological markers have recently become available which reflect conditions of the central nervous system, relating to cell death, axon growth/re-induction, inflammation and/or blood-brain barrier dysfunction. The microtubule-associated protein tau exists in different isoforms, of which 4 to 6 are found in adult brain but only 1 isoform is detected in fetal brain. The diversity of the isoforms is generated from a single gene on human chromosome 17 by alternative mRNA splicing (Himmler, 1989; Goedert et al., 1989; Andreadis et al., 1992). The most striking feature of tau protein, as deduced from molecular cloning, is a stretch of 31 or 32 amino acids, occurring in the carboxy-terminal part of the molecule, which can be repeated either 3 or 4 times. Additional diversity is generated through 29 or 58 amino acid-long insertions in the NH₂-terminal part of tau molecules (Goedert et al., 1989). In vivo tau promotes microtubule assembly and stability in the axonal compartment of neurons by interactions involving its microtubule binding domain which is localised in the repeat region of tau (255-381) (Lewis et al., 1988). In normal circumstances adult brain contains 2 - 3 mol phosphate per mole of tau (Selden and Pollard, 1983; Ksiezak-Reding et al., 1992). Phosphorylation of different sites in normal tau as studied in rat and humans is dependent on the developmental state (Lee et al., 1991; Bramblett et al., 1993; Goedert et al., 1993). Tau variants of 60, 64 and 68 kDa arising as a consequence of phosphorylation have been detected in brain areas showing neurofibrillary tangles (Delacourte et al., 1990; Goedert et al., 1992; Flament and Delacourte, 1990, Greenberg and Davies, 1990). These brains contain 6-8 mol phosphate per mol tau (Ksiezak-Reding et al., 1992). In tau isolated from paired helical filaments, phosphorylation can occur at several positions (Iqbal et al., 1989; Lee et al., 1991; Hasegawa et al., 1992). Detection of normally and abnormally phosphorylated tau in brain extracts is done either via antibodies (Mab Alz50: Ghanbari et al., 1990; Mab Ab423: Harrington et al., 1991; Mab AT120 : Vandermeeren et al., 1993; Mab AT180; Mab AT270 : International application published under WO 95/17429 and Mab AT8 : International application published under WO 93/08302), or via the change in molecular weight (Flament and Delacourte, 1990), or else by functional assay (Bramblett et al., 1992). A combination of monoclonal antibodies, each recognising specific epitopes of tau, has been used to detect the presence of normally and abnormally phosphorylated tau in CSF (Van de Voorde et al., 1995). Tau has been used as a marker to discriminate dementia with altered cytoskeletal properties such as Alzheimer's disease from normal aged subjects or from patients with other types of dementia.

AIMS OF THE INVENTION

It is an aim of the present invention to provide a method for the early detection and/or quantification of CNS damage in an individual, said CNS damage being caused by space-occupying lesions of the CNS, by invasion or metastasis of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents, or by a combination of these mechanisms.

It is a more specific aim of the present invention to provide a method for the early detection and/or quantification of CNS damage in an individual, said CNS damage being caused a primary brain tumour, benign or malignant, brain metastasis or a subdural haematoma.

It is another more specific aim of the present invention to provide a method for the early detection and/or quantification of CNS damage in an individual, said CNS damage being caused by invasion of the CNS by leukemia, lymphoma or breast cancer.

It is another more specific aim of the present invention to provide a method for the early detection and/or quantification of CNS damage in an individual, said CNS damage being caused by bacteria or viruses causing encephalitis or meningitis.

It is another more specific aim of the present invention to provide a method for the early detection and/or quantification of CNS damage in an individual, said CNS damage being caused by stroke, by cerebral infarction, by cerebral haemorrhage, by thrombosis, by perinatal asphyxia, by Binswanger disease or by vasculitis.

It is another more specific aim of the present invention to provide a method for the early detection and/or quantification of CNS damage in an individual, said CNS damage being caused by chemotherapy.

It is another more specific aim of the present invention to provide a method for the early detection and/or quantification of CNS damage in an individual, said CNS damage being caused by trauma, stroke, intracranial pressure or radiation.

It is another aim of the present invention to provide a method for the early detection and/or quantification of CNS damage in an individual, said CNS damage being caused by space-occupying lesions of the CNS, by invasion or metastasis of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents, or by a combination of these mechanisms in order to evaluate the effect of treatment on said CNS damage.

It is another aim of the present invention to provide a kit for the early diagnosis of CNS damage in an individual, said CNS damage being caused by space-occupying lesions of the CNS, by invasion of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents, or by a combination of these mechanisms.

- 5 It is another more specific aim of the present invention to provide a kit for the early diagnosis of CNS damage in an individual, said CNS damage being caused a primary brain tumour, benign or malignant, brain metastasis or a subdural haematoma.

It is another more specific aim of the present invention to provide a kit for the early diagnosis of CNS damage in an individual, said CNS damage being caused by invasion of the CNS by
10 leukemia, lymphoma or breast cancer.

It is another more specific aim of the present invention to provide a kit for the early diagnosis of CNS damage in an individual, said CNS damage being caused by bacteria or viruses causing encephalitis or meningitis.

- 15 It is another more specific aim of the present invention to provide a kit for the early diagnosis of CNS damage in an individual, said CNS damage being caused by stroke, by cerebral infarction, by cerebral haemorrhage, by thrombosis, by perinatal asphyxia, by Binswanger disease or by vasculitis.

It is another more specific aim of the present invention to provide a kit for the early diagnosis of CNS damage in an individual, said CNS damage being caused by chemotherapy.

- 20 It is another more specific aim of the present invention to provide a kit for the early diagnosis of CNS damage in an individual, said CNS damage being caused by trauma, stroke, intracranial pressure or radiation.

It is another aim of the present invention to provide a method to screen or monitor the effect of compounds which prevent or treat CNS damage.

- 25 All the aims of the present invention are considered to be met by the embodiments as set out below.

DETAILED DESCRIPTION OF THE INVENTION

30

The present invention relates to a method for the early detection and/or quantification of CNS

damage in an individual, said CNS damage being caused by space-occupying lesions of the CNS, by invasion of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents or by a combination of these mechanisms. This method comprises the step of determining and/or quantifying the level of tau in an individual and comparing it to the level of tau in control healthy individuals.

The present invention relates to the surprising finding that tau levels in CSF samples from children with leukemia are increased compared to upper limit values for healthy individuals. These increased tau levels are an indication of latent central nervous system invasion and already ongoing CNS damage long before this CNS damage can be measured by the current diagnostic procedures. Also in individuals that were suffering space occupying lesions of the CNS, invasion or metastasis of the CNS, ischemia, stroke or meningitis, increased tau levels were observed in an early stage. Accordingly, tau can be used as an aspecific marker for the early detection of CNS damage caused by invasion of the CNS by leukemia and in general, as an aspecific marker for the early detection of CNS damage caused by CNS damaging agents such as space-occupying lesion of the CNS, invasion or metastasis of the CNS, organisms, anoxia or ischemia, chemical agents, physical agents, or a combination of these mechanisms.

The central nervous system (CNS) is that part of the nervous system which, in vertebrates, consists of the brain and spinal cord, to which sensory impulses are transmitted and from which motor impulses pass out, and which supervises and co-ordinates the activity of the entire nervous system.

The term "CNS damage" refers to any condition of the CNS which is associated with a neuronal malfunctioning and which is caused by a specific inducing agent or damaging agent. More specifically, CNS damage refers to disease processes that include but are not limited to space-occupying lesions of the CNS, invasion or metastasis of the CNS and/or organisms. Space occupying lesions may be, for example, primary brain tumours, benign or malignant, brain metastasis, parasite derived cysts such as *Taenia solium* or *Echinococcus granulosus*, hydrocephalus and/or subdural haematoma. Invasion or metastasis of the CNS can be caused by malignancies such as leukemia, lymphoma, breast cancer, lung cancer, melanoma and/or gastrointestinal malignancies or other types of cancers. Organisms that may infect the CNS and cause CNS damage include but are not limited to prions, viruses, bacteria or parasites. Bacteria and viruses that infect the CNS may cause meningitis, encephalitis, neuro aids or neuroborreliose.

They include but are not limited to *Neisseria meningitidis*, *H. influenzae*, *S. pneumoniae*, *Herpes simplex meningoencephalitis* and *Herpes simplex gluteolis*. CNS damage can also be caused by anoxia or ischemia during anesthesia, perinatal asphyxia, drowning, asthma, stroke, cerebral infarction, thrombosis, cerebral haemorrhage, CO poisoning, Binswanger disease and/or vasculitis.

5 Chemical agents causing CNS damage include but are not limited to gene therapy, pharmaceuticals, chemotherapy and exposure to chemical compounds. CNS damage can also be caused by physical agents such as trauma, radiation, hypothermia, hyperthermia, intracranial pressure or stroke. It is also possible that more than one of the above mentioned causing agents are responsible for the CNS damage.

10 The present invention thus provides a method for the early detection and/or quantification of said CNS damage by determining the level of tau. "Early detection and/or quantification of CNS damage" means that the CNS damage is determined by a method that allows it to be detected before it is detectable by the current methods.

The term "tau" as referred to in the present application can be any form of tau, including any state of phosphorylation. The level of tau is determined qualitatively and/or quantitatively as a measure for the degree of CNS damage. Tau can be detected in vitro as well as in vivo.

15 The method for the early in vitro detection of CNS damage in an individual comprises the steps of obtaining a sample from said individual, determining and/or quantifying the level of tau in said sample and comparing it to the level of tau in a sample of control healthy individuals. The term "sample" refers to any source of biological material, for instance body fluids, hair, epithelial cells, peripheral blood or any other sample comprising tau protein. In a preferred embodiment, tau can be detected and/or quantified in vitro by analysis of the level of tau in a body fluid sample of the patient. The term "body fluid" refers to all fluids that are present in the human body including but not limited to blood, lymph, urine and cerebrospinal fluid (CSF). In a more specific embodiment
20 of the present invention tau is detected and/or quantified in a cerebrospinal fluid sample taken from the patient. In another specific embodiment of the invention tau is detected and/or quantified in a sample of blood derivatives of the patient. The blood sample can include the whole sample as taken from the patient. More preferably the blood sample includes a plasma sample or a serum sample.

30 Tau can be detected and/or quantified by any method known, including but not limited to the use of antibodies, the change in molecular weight (Flament and Delacourte, 1990), or else by

functional assay (Bramblett et al., 1992). In a preferred embodiment tau can be detected by an immunoassay comprising at least the following steps:

- obtaining a sample from the patient; and
- bringing said sample into contact with a monoclonal antibody (primary antibody or capturing antibody) recognising tau, under conditions being suitable for producing an antigen-antibody complex; and
- detecting the immunological binding of said antibody to said sample.

Advantageously, the monoclonal antibody used in the invention is in an immobilised state on a suitable support. Alternatively, the present process may be put into practice by using any other immunoassay format known to the person skilled in the art.

The process for the detection of the antigen can then be carried out by bringing together said antigen-antibody complex formed by the antigen and the primary antibody recognising tau with:

- a) a secondary antibody (or detector antibody)

*which can be a monoclonal antibody recognising an epitope of the tau-primary antibody complex but not recognising the primary antibody alone, or

*which can be a polyclonal antibody recognising an epitope of the tau-primary antibody complex but not recognising the primary antibody alone, with said polyclonal antibody being preferably purified by immunoaffinity chromatography using immobilised tau or the tau-primary antibody complex.

- b) a marker either for specific tagging or coupling with said secondary antibody, with said marker being any possible marker known to the person skilled in the art;
- c) appropriate buffer solutions for carrying out the immunological reaction between the primary antibody and the sample, between the secondary antibody and the tau-primary antibody complex and/or between the bound secondary antibody and the marker, and,
- d) possibly also, for standardisation purposes, a purified protein or synthetic peptide reactive with the antibodies that recognise tau.

Advantageously, the secondary antibody itself carries a marker or a group for direct or indirect coupling with a marker.

The term "epitope" refers to that portion of the antigen-antibody complex that is specifically bound by an antibody-combining site. Epitopes may be determined by any of the techniques known in the art or may be predicted by a variety of computer prediction models known in the

art.

The expression “recognising”, “reacting with”, “immunological binding” or “producing an antigen-antibody complex” as used in the present invention is to be interpreted that binding between the antigen and antibody occurs under all conditions that respect the immunological properties of the antibody and the antigen.

Any monoclonal or polyclonal antibody that specifically recognises tau may be used for the detection of tau. Antibodies specifically recognising normally and/or abnormally phosphorylated tau include Alz50 (Ghanbari et al., 1990), Ab423 (Harrington et al., 1991), AT8 (International application published under WO 93/08302), AT120 (Vandermeeren et al., 1993), AT180 and AT270 (International application published under WO 95/17429) and AT100 (International application published under WO 96/04309). But also other antibodies known in the art that specifically recognise tau can be used.

The method for the early in vitro detection and/or quantification of CNS damage in an individual can also be used to evaluate the effect of a certain treatment on the CNS damage in said individual. Possible treatments that might influence the status of the CNS include but are not limited to drug treatments, chemotherapy, physical therapy, including radiotherapy and gene therapy.

The method for the early in vivo detection and/or quantification of CNS damage in an individual comprises the steps of determining and/or quantifying the level of tau in said individual and comparing it to the level of tau in control healthy individuals. In a preferred embodiment, tau can be detected in vivo by in vivo imaging. Tau can be visualised in situ by non-invasive methods including but not limited to brain imaging methods described by Arbit et al. (1995), Tamada et al. (1995), Wakabayashi et al. (1995), Huang et al. (1996), Sandrock et al. (1996), Mariani et al. (1997). These in vivo imaging methods may allow the localisation and visualisation of tau, for example, by use of labelled antibodies recognising tau.

Tau can also be used as an aspecific marker for in vivo imaging to evaluate the effect of a certain treatment on the CNS damage in an individual. Possible treatments that might influence the status of the CNS include but are not limited to drug treatments, chemotherapy, physical therapy, including radiotherapy and gene therapy.

The present invention further relates to the use of tau as an aspecific marker for the manufacture of a diagnostic kit for the early detection in an individual of CNS damage caused by space-

occupying lesions of the CNS, by invasion or metastasis of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents, or by a combination of these mechanisms

The present invention further relates to the use of tau as an aspecific marker for the manufacture of a diagnostic kit for the early detection in an individual of CNS damage caused by a primary

5 brain tumour, benign or malignant, brain metastasis, or a subdural haematoma.

The present invention further relates to the use of tau as an aspecific marker for the manufacture of a diagnostic kit for the early detection in an individual of CNS damage caused by invasion or metastasis of the CNS by leukemia, lymphoma or breast cancer.

10 The present invention further relates to the use of tau as an aspecific marker for the manufacture of a diagnostic kit for the early detection in an individual of CNS damage caused by bacteria or viruses.

15 The present invention further relates to the use of tau as an aspecific marker for the manufacture of a diagnostic kit for the early detection in an individual of CNS damage caused by stroke, by cerebral infarction, by thrombosis, by cerebral haemorrhage, by perinatal asphyxia, by Binswanger disease or by vasculitis.

The present invention further relates to the use of tau as an aspecific marker for the manufacture of a diagnostic kit for the early detection in an individual of CNS damage caused by chemotherapy.

20 The present invention further relates to the use of tau as an aspecific marker for the manufacture of a diagnostic kit for the early detection in an individual of CNS damage caused by trauma, stroke, intracranial pressure or radiation.

25 The present invention further relates to a kit for the in vitro or in vivo diagnosis in an individual of CNS damage caused by space-occupying lesions of the CNS, by invasion or metastasis of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents, or by a combination of these mechanisms. Any kit that provides a tool for the detection of tau can be used for the diagnosis of the above-mentioned CNS damage.

30 A preferred kit for the in vitro diagnosis in an individual of CNS damage caused by space-occupying lesions of the CNS, by invasion or metastasis of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents, or by a combination of these mechanisms is based on an immunoassay and comprises:

- at least a monoclonal antibody (primary antibody) which forms an immunological

complex with an epitope of the tau protein;

- a secondary antibody

* which can be a monoclonal antibody recognising an epitope of the tau-primary antibody complex but not recognising the primary antibody alone, or

5 * which can be a polyclonal antibody recognising an epitope of the tau-primary antibody complex but not recognising the primary antibody alone, with said polyclonal antibody being preferably purified by immunoaffinity chromatography using immobilised tau protein or immobilised tau-primary antibody complex;

10 - a marker either for specific tagging or coupling with said secondary antibody;

- appropriate buffer solutions for carrying out the immunological reaction between the primary antibody and a test sample, between the secondary antibody and the tau-primary antibody complex, and/or between the bound secondary antibody and the marker;

15 - possibly, for standardisation purposes, a purified protein or synthetic peptide containing one of more tau epitopes.

The present invention also relates to a method to screen or monitor the effect of compounds which prevent or treat CNS damage comprising the step of determining the level of tau and comparing it to the level of tau in a control sample.

20 The present invention will now be illustrated by reference to the following examples that set forth particularly advantageous embodiments. However, it should be noted that these examples are illustrative and can not be construed as to restrict the invention in any way.

Table 1: Treatment Protocol for patients with non-B-cell ALL/NHL (EORTC 58881).

| Treatment phase | Drugs | Dose | Route | Days |
|--|---------------------|-------------------------|-------|---|
| Prephase | Prednisolone | 60 mg/m ² /d | PO | day 1 - 7 |
| | MTX | according to age | IT | day 1 (<u>LP1</u>), day 8 (<u>LP2</u>), day 22 (<u>LP3</u>) |
| Protocol I: induction (37 days) | Prednisolone | 60 mg/m ² /d | PO | day 8 - 28 |
| | | 20 mg/m ² /d | PO | day 29 - 31 |
| | | 10 mg/m ² /d | PO | day 32 - 34 |
| | | 5 g/m ² /d | PO | day 35 - 37 |
| | Vincristine | 1.5 mg/m ² | IV | day 8, 15, 22, 29 |
| | Daunorubicine | 30 mg/m ² | IV | day 8, 15, 22, 29 |
| | E coli asparaginase | 10000 U/m ² | IV | day 12, 15, 19, 22, 25, 29, 32, 35 |
| Protocol I: consolidation (26 days) | Cyclophosphamide | 1000 mg/m ² | IV | day 36, 63 |
| | 6-mercaptopurine | 60 mg/m ² | PO | day 36 - 63 |
| | Ara-C | 75 mg/m ² | IV | day 38 - 41, 45 - 48, 52 - 55, 59 - 62 |
| | MTX | according to age | IT | day 38 (<u>LP4</u>), 52 (<u>LP5</u>) |
| (14 days) | | | | |

Table 1. Continued.

| Treatment phase | Drugs | Dose | Route | Days |
|---|---------------------|-----------------------------|-------|---|
| Interval therapy (56 days) | 6-mercaptopurine | 25 mg/m ² | PO | day 1 - 56 |
| | MTX | 5000 mg/m ² | IV | day 8, 22, 36, 50 |
| | MTX | according to age | IT | day 9 (<u>LP6</u>), 23 (<u>LP7</u>), 37 (<u>LP8</u>), 51 (<u>LP9</u>) |
| (14 days) | | | | |
| Protocol II: induction (35 days) | Dexamethasone | 6 mg/m ² /d | PO | day 1 - 21 |
| | | 3 mg/m ² /d | PO | day 22 - 35 |
| | | 1 mg/m ² /d | PO | day 26 - 29 |
| | Vincristine | 1.5 mg/m ² | IV | day 8, 15, 22, 29 |
| | Adriamycine | 30 mg/m ² | IV | day 8, 15, 22, 29 |
| | E coli asparaginase | 10000 U/m ² | IV | day 8, 11, 15, 18 |
| Protocol II: consolidation (14 days) | Cyclophosphamide | 1000 mg/m ² | IV | day 36 |
| | 6-thioguanine | 60 mg/m ² | PO | day 36 - 49 |
| | Ara-C | 75 mg/m ² | IV | day 38 - 41, 45 - 48 |
| | MTX | according to age | IT | day 38 (<u>LP10</u>) |
| | | | | |
| (14 days) | | | | |
| Maintenance | 6-mercaptopurine | 50 mg/m ² /d | PO | Total duration of treatment = 2 years |
| | MTX | 20 mg/m ² weekly | PO | |

Table 2. Treatment protocol for patients with B-cell NHL (UKCCSG 9602).

| Treatment phase | Drugs | Dose | Route | Days |
|--|------------------|------------------|-------|--|
| COP (7 Days) | Prednisolone | 60 mg/m2/d | PO | day 1 – 7 |
| | Vincristine | 1 mg/m2 | IV | day 1 |
| | Cyclophosphamide | 300 mg/m2 | IV | day 1 |
| | MTX | according to age | IT | day 1 (<u>LP1</u>) |
| | Hydrocortisone | according to age | IT | day 1 (<u>LP1</u>) |
| *** Start depending on the status of the patient | | | | |
| COPADM 1 and COPADM 2 (5 days) | Prednisolone | 60 mg/m2/d | PO | day 1 – 5 |
| | Vincristine | 2 mg/m2 | IV | day 1 |
| | Cyclophosphamide | 500 mg/m2/d | IV | day 2 – 4 |
| | Doxorubicine | 60 mg/m2 | IV | day 2 |
| | MTX | according to age | IT | day 2 (<u>LP2</u>), 6 (<u>LP3</u>) |
| | Hydrocortisone | 3000 mg/m2 | IV | day 1 |
| | | according to age | IT | day 2 (<u>LP2</u>), 6 (<u>LP3</u>) |
| *** Start depending on the status of the patient | | | | |

Table 2. Continued.

| Treatment phase | Drugs | Dose | Route | Days |
|--|------------------|--------------------------|-------|--|
| CYM 1 and CYM 2 (6 days) | MTX | according to age | IT | day 2 (<u>LP4</u>) |
| | | 3000 mg/m ² | IV | day 1 |
| | Hydrocortisone | according to age | IT | day 2 (<u>LP4</u>), 7 (<u>LP5</u>) |
| | Ara-C | according to age | IT | day 7 (<u>LP5</u>) |
| | | 100 mg/m ² /d | IV | day 2-6 |
| *** Start depending on the status of the patient | | | | |
| COPADM 3 (5 days) | Prednisolone | 60 mg/m ² /d | PO | day 1 - 5 |
| | Vincristine | 2 mg/m ² | IV | day 1 |
| | Cyclophosphamide | 500 mg/m ² /d | IV | day 2 - 3 |
| | Doxorubicine | 60 mg/m ² | IV | day 2 |
| | MTX | according to age | IT | day 2 |
| | | 3000 mg/m ² | IV | day 1 |
| | Hydrocortisone | according to age | IT | day 2 |

IT = intra thecal; IV = intravenous, PO = per os, LP = lumbar puncture, MTX = methotrexate, Ara-C = arabinoside

Table 3: Treatment Protocol for patients with AML (EORTC 58921).

| Treatment phase | Drugs | Dose | Route | Days |
|------------------------|---------------|---------------------------|-------|--|
| Induction | Ara-C | 100 mg/m ² | IV | day 1, 2 |
| | Ara-C | 200 mg/m ² /d | IV | day 3 - 8 |
| | Mitoxantrone | 10 mg/m ² /d | IV | day 3 - 5 |
| | VP16 | 150 mg/m ² /d | IV | day 6 - 8 |
| First intensification | Ara-C | according to age | IT | day 1 (<u>LP1</u>), 8 (<u>LP2</u>) |
| | Ara-C | 3000 mg/m ² /d | IV | day 1 - 4 |
| | Mitoxantrone | 10 mg/m ² /d | IV | day 5 - 7 |
| Second intensification | Daunorubicin | 20 mg/m ² /d | IV | day 1 - 4 |
| | Ara-C | 200 mg/m ² /d | IV | day 1 - 4 |
| | VP16 | 100 mg/m ² /d | IV | day 1 - 4 |
| | 6-thioguanine | 100 mg/m ² /d | PO | day 1 - 4 |
| | Dexamethasone | 6 mg/m ² /d | PO | day 1 - 4 |
| | Ara-C | according to age | IT | day 1 (<u>LP3</u>), 4 (<u>LP4</u>) |
| Third intensification | Ara-C | 2000 mg/m ² /d | IV | day 1 - 3 |
| | VP16 | 125 mg/m ² /d | IV | day 2 - 5 |
| Maintenance | 6-thioguanine | 40 mg/m ² | PO | 1 year |
| | Ara-C | 40 mg/m ² | SC | 4 days/month |

IT = intra thecal; IV = intravenous, PO = per os, SC = subcutaneous, LP = lumbar puncture, Ara-C = arabinoside, VP16 =

Table 4. Level of tau in CSF samples of patients with possible CNS damage caused by different factors.

| Cause of CNS damage ^c | Centre ^a | Patient | Age | Sex ^b | Tau level (pg/ml) |
|--|---------------------|---------|-----|------------------|----------------------|
| SPACE OCCUPYING LESIONS, INVASION OR METASTASIS | | | | | |
| Subdural haematome | 01 | 025 | 39 | M | 263 |
| Anaplastic oligoastrocytoma | 01 | 032 | 68 | M | 329 |
| Cerebral metastasis | 08 | 024 | 70 | F | 292 |
| Oligodendroglioma WHO III | 08 | 031 | 48 | M | 335 |
| Metastasis of breast cancer | 08 | 020 | 52 | F | 150 |
| BLEEDING, INFARCTION OR ISCHEMIA | | | | | |
| Stroke - CVA | 01 | 021 | 62 | F | 451 |
| Multiple lacunar stroke | 01 | 026 | 56 | F | 250 |
| Stroke - CVA | 04 | 004 | 50 | F | 105 |
| Acute but limited stroke | 05 | 032 | 72 | M | 248 |
| Subarachnoid hemorrhage | 08 | 019 | 50 | M | 216 |
| Chronic post-anoxia (vegetative state) | 05 | 051 | 37 | F | 446 |
| Binswanger disease | 10 | 014 | 74 | M | 347 |
| Vasculitis | 10 | 009 | 57 | F | 1250 |
| Cerebral ischemia | 10 | 005 | 80 | M | 273 |
| Ischemia | 10 | 006 | 71 | F | 574 |
| Superior sagittal sinus thrombosis | 01 | 023 | 41 | M | 773 |

Table 4. Continued.

| Cause of CNS damage ^c | Centre ^a | Patient | Age | Sex ^b | Tau level (pg/ml) |
|---|---------------------|---------|-----|------------------|----------------------|
| ORGANISMS | | | | | |
| Cryptococcal meningitis | 05 | 033 | 29 | M | 1143 |
| Neuroborreliose | 05 | 045 | 16 | M | 141 |
| Chronic meningitis, unknown cause | 05 | 046 | 34 | F | 452 |
| Meningococcal meningitis | 05 | 049 | 24 | M | 37 |
| Bacterial meningitis | 05 | 059 | 73 | F | 446 |
| Neuro aids | 05 | 060 | 45 | F | 237 |
| Pneumococcal meningitis | 05 | 063 | 70 | M | 111 |
| Bacterial meningitis | 06 | 026 | 25 | M | 1250 |
| Bacterial meningitis | 06 | 027 | 19 | M | 37 |
| TBC meningitis | 06 | 028 | 54 | M | 720 |
| Bacterial meningitis | 06 | 029 | 23 | M | 37 |
| Bacterial meningitis | 06 | 030 | 28 | M | 88 |
| Viral meningitis | 06 | 037 | 82 | M | 314 |
| Bacterial meningitis | 08 | 018 | 49 | F | 205 |
| Viral meningitis | 08 | 029 | 29 | F | 170 |
| CONTROLS | | | | | |
| Muscle weakness and hysterical conversion | 01 | 020 | 39 | F | 282 |
| Facial palsy (peripheral nerve disease, normal CSF | 05 | 041 | 44 | M | 141 |
| Tension (psychogenic) headache | 05 | 031 | 49 | F | 37 |
| Psychogenic headache and neurosis | 05 | 036 | 56 | F | 131 |
| Psychogenic signs and symptoms | 05 | 040 | 48 | F | 220 |
| Major hysteria | 05 | 042 | 47 | F | 471 |
| Lumbar disc prolaps and sciatica | 05 | 047 | 74 | M | 195 |

Table 4. Continued.

| Cause of CNS damage ^c | Centre ^a | Patient | Age | Sex ^b | Tau level (pg/ml) |
|--------------------------------------|---------------------|---------|-----|------------------|----------------------|
| CONTROLS | | | | | |
| Control (associated Sjogren disease) | 05 | 065 | 52 | F | 107 |
| Control (neurosis) | 05 | 079 | 40 | F | 90 |
| Depression | 10 | 008 | 59 | F | 201 |
| Depression | 10 | 010 | 69 | F | 139 |
| Myalgia, myositis | 04 | 003 | 31 | M | 37 |
| Myalgia, myositis | 04 | 005 | 35 | F | 435 |
| Malaise, fatigue | 04 | 006 | 50 | M | 98 |
| Neck pain, vertigo | 08 | 011 | 39 | F | 192 |
| Headache | 08 | 017 | 49 | M | 113 |
| Depression | 08 | 021 | 57 | M | 361 |
| Headache, strabismus convergens | 08 | 033 | 32 | F | 239 |
| Bell's palsy | 06 | 031 | 38 | M | 90 |
| Bell's palsy | 06 | 032 | 22 | F | 131 |
| Bell's palsy | 06 | 033 | 73 | F | 231 |
| Bell's palsy | 06 | 034 | 74 | F | 144 |
| Carpal tunnel syndrome | 06 | 035 | 71 | M | 190 |
| Carpal tunnel syndrome | 06 | 036 | 90 | M | 665 |

^a 01: Prof. P. Cras, University Hospital, Neurology, Antwerp, Belgium and Dr. A. Daniels, UIA, Laboratory of Neurobiology, Antwerp, Belgium; 04: Dr. P.D. Mehta, Institute Basic Research, Staten Island, NY, USA; 05: Dr. A. Ivanoiu, UCL St-Luc, Laboratory of Neurochemistry, Brussels, Belgium; 06: Prof. P.P. De Deyn, UIA, Laboratory of Neurochemistry, Antwerp, Belgium; 08: Dr. C. Bancher, Lainz Hospital, Neurology, Vienna, Austria; 10: Dr. J. Wiltfang, Georg-August University, Gerontopsychiatry, Göttingen, Germany.

^b F: female; M: male.

^c In cases where multiple factors may contribute to the damage, only the factor supposed to be

most relevant is given.

5 FIGURE LEGENDS

Figure 1. Tau values at diagnosis, before any treatment was given: 1. Control children; 2. AML; 3. AML-CNS+; 4. CML; 5. MDS; 6. B-NHL; 7. Non-B-ALL; 8. Non-B-ALL CNS+; 9. VHR non-B-ALL.

Figure 2. CSF-tau levels in seven patients after acute ischemic stroke. The CSF samples were collected at income (day 0-1), day 2-3, day 7-8, day 21-22 (3 weeks) and day 90-110 (3 months).

Figure 3. CSF-tau levels at the time of maximal release in relation to the size of the infarction as measured by CT scan.

EXAMPLES

Example 1: Increased tau levels in children with leukemia

To evaluate the influence of chemotherapy on neuronal damage, a longitudinal study was conducted, involving 65 children with leukemia (aged 2 to 16 years) without measurable central nervous system involvement, and treated according to standard procedures. A total of 377 CSF samples were analysed. Before each injection, a small volume of fluid was sampled for routine laboratory analysis and the leftover was used in our study. These children were being diagnosed and then treated for their leukemia at the University Hospital of Leuven, Belgium. Tau protein in cerebrospinal fluid was assessed using the INNOTEST hTAU Antigen (Innogenetics, Gent, Belgium).

For all children suspected to have leukemia, a lumbar puncture was performed before the start of

the treatment to detect the possible presence of leukemic cells, indicative of central nervous system invasion. The tau levels measured at that time, and thus before treatment, would serve as the control level to compare chemotherapy-induced changes in the levels of tau.

We observed that some children with leukemia already had very high tau levels at diagnosis in spite of the fact that no leukemic cells were detected in the central nervous system. These children constitute a new risk group having brain invasion or leukemia-induced CNS damage, which cannot always be found using current diagnostic procedures (imaging of the brain, lumbar puncture, eye funduscopy). This was further supported by the increased tau levels seen in one patient having leukemia with proven cellular invasion into the brain (malignant cells in the cerebrospinal fluid).

Example 2: Increased tau levels in patients with leukemia before treatment

1. Subjects

Between August 1996 and June 1999, 510 samples of CSF were taken from 82 children being treated for cancer at the Pediatric Hemato-oncology Department of the Catholic University of Leuven, Belgium. CSF samples were only taken in the course of scheduled lumbar punctions (LPs) for staging or treatment for malignancy.

Three groups of patients with hematological malignancies were studied. The largest group consisted of 48 patients with non-B-ALL, treated according to the EORTC protocol 58881 (table 1). Of these children, 20 had CD10(+) blasts (or common ALL), from whom two patients had also Down syndrome (DS) and 1 patient had the Brachmann-de Lange syndrome; 6 patients had common B-cell blasts, 2 patients had common T-cell blasts, 2 patients had pro-B-cell blasts, 9 patients had pre-B-cell blasts, and 9 patients had T-cell blasts. Forty-two children had leukemia, 6 patients had non-Hodgkin's lymphoma stage II (1 patient), Stage III (4 patients) or Stage IV (1 patient). One patient had overt CNS involvement (CNS+), defined according to the study protocol with malignant cells in the CSF. Five patients were considered as very high risk (VHR) patients according to the criteria defined in the protocol (2 patients with t(9;22) and 3 patients

with corticoid-resistance). As the first part of the induction treatment was similar to the other patients, they were included in the analyses according to the induction chemotherapy. Twenty-eight of the patients within non-B-ALL could be followed longitudinally during their treatment period.

- 5 A second group of patients included 10 B-cell non-Hodgkin's lymphoma patients, treated according to the United Kingdom Children Cancers group (UKCCSG 9602) NHL protocol (table 2). Five patients had B-cell lymphomas, 3 patients had Burkitt's lymphoma, and 2 patients had anaplastic large cell lymphoma (ALCL). All patients were treated with the same protocol, except one patient who had B-cell leukemia. Six of these patients were studied longitudinally.
- 10 A third patient group consisted of 9 children with Acute Myeloid Leukemia - myelodysplasia (AML - MDS), of which 2 patients had CNS involvement and two patients had Down syndrome. There were 3 patients with M0, and 1 patient each with M1, M2, M5a or M7 phenotype. Two patients had MDS, of which one had developed AML and was treated with chemotherapy. All these patients, except one MDS patient, were treated according to the EORTC 58921 protocol
- 15 (table 3) and 7 patients were followed longitudinally.

- The other patients consisted of a heterogeneous group of children ($n = 9$) in which for clinical reasons an LP was performed. This group includes 3 children with medulloblastoma (staging), 2 children with rhabdomyosarcoma (staging), 1 child with Langerhans cell histiocytosis (LCH, staging), ganglioglioma (staging), germinoma (staging), and retinoblastoma with CNS metastasis
- 20 (staging and follow up). The control group consisted of 4 children from whom CSF was taken as part of the routine control for possible viral or bacterial infections, but with negative results. One patient with a localized retinoblastoma (staging) and one screened for familial hemophagocytic lymphohistiocytosis (HLH) were also included in the control group. The nearest relatives of the patients gave oral informed consent for participation in the study.

25

2. Methods

- Liquor sampling.** Lumbar punctures were performed just prior to the IT administration of therapeutics. Five ml of CSF was collected in different polypropylene tubes. One sample was
- 30 centrifuged immediately at 1500 rpm for 2 minutes to eliminate cells and other insoluble material. The supernatant was stored at -70°C for subsequent analysis. The number of freeze/thaw cycles

was restricted to a minimum.

Measurement of tau in the CSF. All biochemical analyses for the detection of CSF-tau were made without knowledge of the clinical diagnosis. Potentially confounding factors, such as the number of freeze/thaw cycles, recipient type, and volume of sample per assay were standardized for the whole study protocol. CSF-tau levels were determined using a sandwich ELISA (INNOTEST hTAU Antigen, Innogenetics, Gent, Belgium), that measured total tau (both normal and hyperphosphorylated tau). Samples were analyzed firstly alone, at the time of each LP. Afterwards, all samples derived from one patient were analyzed again on one immunoplate. The correlation coefficient between the results from the first and the second approach for a set of 104 samples was 0.901 (95%CI: 0.856 - 0.933). The increases of CSF-tau described in the paper were excluded to be general increases of proteins in the liquor.

3. Results

Normal upper limit levels for tau were firstly determined on CSF samples from the 6 control children. The mean CSF-tau value was 106.2 pg/ml (95%CI = 34.3-178.0). Arbitrary cut-off normal value was considered as 312 pg/ml (mean + 3 SD), which is in the range of values observed in adults (Hulstaert et al., 1999). Moreover, we did not find a correlation between the CSF-tau values at diagnosis and the age of the children (Pearson $r = -0.161$, $CI_{95\%} = -0.3914 - 0.08836$, $n = 64$), suggesting that age has no influence on CSF-tau levels. Clearly pathological CSF-tau values can be considered as above 500 pg/ml.

Tau levels at diagnosis were analyzed for each subgroup of patients (Fig. 1). There was no obvious difference for the tau levels in patients with and without Dow's syndrome (not shown). The two patients with overt CNS invasion (CNS+) had levels of CSF-tau at diagnosis above 312 pg/ml. However, the 2 children with MDS, 7/28 children with non-B-ALL, 1/4 non-B-ALL patients with very high risk criteria, 1/5 patients with AML, and 2/8 patients with B-cell NHL had a level of tau above 312 pg/ml, while using classical diagnostic procedures, CNS invasion was not detected. Three patients with risen intracranial pressure (medulloblastoma), and one patient with germinoma, from which CSF was taken for staging, had high CSF-tau concentrations

(823,1397,1500 and 442 pg/ml), in contrast to one patient with grade I astrocytoma who had a normal CSF-tau level (97 pg/ml). One patient with LCH had a normal CSF-tau level of 112 pg/ml. Two patients with rhabdomyosarcoma could be analyzed at diagnosis: one patient with stage I disease had a CSF-tau level of 279 pg/ml, the other patient with stage IV disease had a level of 320 pg/ml. One patient with retinoblastoma and CNS involvement had a CSF-tau level of 1800 pg/ml.

For 33 patients with non-B-ALL, CSF-tau levels at diagnosis did not correlate with tumor burden, as reflected by the white blood cell count (Pearson $r = 0.04575$, CD95%: -0.3024 - 0.3831) or serum LDH (Pearson $r = -0.03002$, CI95%: -0.3696 - 0.3166). In patients with B-NHL, there was no significant correlation between the LDH level and CSF-tau (Pearson $r = -0.3723$, CD95%: -0.8532 - 0.4507).

Example 3: Use CSF-tau as a marker for early detection of possible CNS damage caused by stroke

1. Subject

Seven patients, 3 men and 4 women, 63-81 years old (mean SD, 70.7 ± 7.2 years) with cerebral infarctions admitted to the Unit of Neurology, Sahlgren's University Hospital, Göteborg, Sweden, were incorporated in the study. All patients were included within 72 h of stroke onset.

2. Methods

CSF samples were collected using lumbar puncture. 12 ml was collected and frozen in 0.5 ml aliquots at -80°C until analyzed. CSF samples were collected at income (day 0-1), day 2-3, day 7-8, day 21-22 (3 weeks) and day 90-110 (3 months). CSF-tau levels were determined using a sandwich ELISA (INNOTEST hTAU Antigen, Innogenetics, Gent, Belgium), that measured total tau (both normal and hyperphosphorylated tau).

The extent of brain damage was examined with computerized tomography (CT) of the brain during the first day of admittance. Clinical the patients were also examined using the modified

Scandinavian Stroke Scale Index (SSI; Scandinavian Stroke Study Group, 1985) at onset of stroke and 3 months later for the degree of disability with the Bartel Index (BI; Mohoney and Barthel, 1965).

5 3. Results

CSF-tau showed a marked increase after acute stroke, with a peak after 1-3 weeks and return to normal after 3 months (Fig 2). There was also a correlation between CSF-tau levels and the size of the infarction as measured by CT scan (Fig 3). These results indicate that CSF-tau reflects neuronal damage and degeneration and the level in the CSF depends on the amount of damaged nerve cells. In this study no correlation was found between the clinical data (SSI or BI) and CSF-tau, probably due to small number of patients.

15 Example 4: Use of tau as a marker for early detection of possible CNS damage caused by different damaging agents

1. Subjects

20 A multicentre study was carried out at 8 European and 2 U.S. university centres involved in CSF research, based on residual CSF archived at the centres for research purposes. CSF samples of patients with a broad range of different neurological conditions were included in order to get a general idea about the specificity of the tau marker changes in a variety of pathologies involving the CNS. Neurological controls consisted of subjects without obvious CNS damage (table 4).

25 The study was conducted in accordance with local clinical research regulations. If required additional local Ethics Committee or Institutional Review Board approval was obtained by the investigator prior to the start of the study.

30 2. Methods

CSF samples were collected using lumbar puncture (LP). Only CSF samples containing less than 500 red blood cells per μl were included in the study. The CSF samples were centrifuged at 2000 g for 10 minutes within 4 hours after LP and kept frozen without thawing. CSF samples from centre 01 had undergone an additional freeze-thaw cycle before analysis. The concentration of total tau comprising normal tau and paired helical filament-tau was measured at the centres using a sandwich ELISA technique (INNOTEST hTAU-Antigen, Innogenetics N.V.).

3. CSF-tau levels in patients with possible CNS damage compared to CSF-tau levels in neurological control subjects

Tau levels in the CSF samples of patients with possible CNS damage caused by space occupying lesions, by invasion or metastasis, by bleeding, infarction or ischemia or by organisms and tau levels in CSF samples of control subjects are shown in table 5. The group of patients that had possibly contracted CNS damage by space occupying lesions, by invasion or metastasis, by bleeding, by infarction or ischemia, or by organisms showed an overall higher tau level than the neurological control patients ($p = 0.022$, two sided Mann Whitney test).

REFERENCES

Andreadis A., Brown W., Kosik K. (1992) Structure and novel exons of the human tau gene. *Biochem.* 31: 10626-10633.

Arbit E., Cheung N.K., Yeh S.D., Daghighian F., Shang J.J., Cordon-Cardo C., Pentlow K., Canete A., Finn R., Larson S.M. (1995) Quantitative studies of monoclonal antibody targeting to disialogangliosid GD2 in human brain tumours. *Eur. J. Nucl. Med.* 22: 419-426.

Bramblett G., Trojanowski J., Lee V. (1992) Regions with abundant neurofibrillary pathology in human brain exhibit a selective reduction in levels of binding-competent tau and accumulation of abnormal tau isoforms (A68 proteins). *Lab. Invest.* 66: 212-222.

Bramblett G., Goedert M., Jakes R., Merrick S., Trojanowski J., Lee V. (1993) The abnormal phosphorylation of tau at Ser396 in Alzheimer's disease recapitulates phosphorylation during development and contributes to reduced microtubule binding. *Neuron*. 10: 1089-1099.

- 5 Delacourte A., Flament S., Dibe E., Hublau P., Sablonniere B., Hemon B., Sherrer V., Defossez A. (1990) Pathological proteins Tau64 and 69 are specifically expressed in the somatodendritic domain of the degenerating cortical neurons during Alzheimer's disease. *Acta Neuropathol*. 80: 111-117.
- 10 Flament S., Delacourte A. (1990) Tau Marker? *Nature* 346: 6279.
- Ghanbari H., Kozuk T., Miller B., Riesing S. (1990) A sandwich enzyme immunoassay for detecting and measuring Alzheimer's disease-associated proteins in human brain tissue. *J. Clin. Laboratory Anal.* 4: 189-192.
- 15 Goedert M., Spillantini M., Jakes R., Rutherford D., Crowther R. (1989) Multiple isoforms of human microtubule-associated protein tau: sequences and localisation in neurofibrillary tangles of Alzheimer's disease. *Neuron*. 3: 519-526.
- 20 Goedert M., Cohen E., Jakes R., Cohen P. (1992) p42 Map kinase phosphorylation sites in microtubule-associated protein tau one dephosphorylated by protein phosphatase 2A1: implications for Alzheimer's disease. *FEBS Lett.* 312: 95-99.
- Goedert M., Jakes R., Crowther R., Six J., Lübke U., Vandermeeren M., Cras P., Trojanowski J.Q., Lee V. (1993) The abnormal phosphorylation of tau protein at serine 202 in Alzheimer's disease recapitulates phosphorylation during development. *Proc. Natl. Acad. Sci. (USA)* 90: 5066-5070.
- 25 Greenberg S., Davies P. (1990) A preparation of Alzheimer paired helical filaments that displays distinct tau proteins by polyacrylamide gel electrophoresis. *Proc. Natl. Acad. Sci. USA* 87: 5827-5831.
- 30

- Harrington C., Mukaetova E., Hills R., Edwards P., Montejo de Garcini E., Novak M., Wischik C. (1991) Measurement of distinct immunochemical presentations of tau protein in Alzheimer's disease. *Proc. Natl. Acad. Sci. (USA)* 88: 5842-5846.
- 5 Hasegawa M., Morishima-Kawashima M., Takio K., Suzuki M., Titani K., Ihara Y. (1992) Protein sequence and mass spectrometric analyses of tau in Alzheimer's disease brain. *J. Biol. Chem* 267: 17047-17054.
- 10 Himmler A. (1989) Structure of the bovine tau gene: alternatively spliced transcripts. *Mol. Cell Biol.* 9(4): 1389-96.
- Huang Q., He G., Lan Q., Li X., Qian Z., Chen J., Lu Z., Du Z. (1996) Target imaging diagnosis of human brain glioma. Clinical analysis of 40 cases. *Nucl. Med. Commun.* 17: 311-316.
- 15 Hulstaert F., Blennow K., Ivanoiu A., Schoonderwaldt H.C., Riemenschneider M., De Deyn P.P., Bancher C., Cras P., Wiltfang J., Mehta P.D., Iqbal K., Pottel H., Vanmechelen E., Vanderstichele H. (1999) Improved discrimination of AD patients using β -amyloid₍₁₋₄₂₎ and tau levels in CSF. *Neurology* 52: 1555-1562.
- 20 Iqbal K., Grundke-Iqbal I., Smith A., George L., Tung Y., Zaidi T. (1989) Identification and localisation of a Tau peptide to paired helical filaments of Alzheimer's disease. *Proc. Natl. Acad. Sci. (USA)* 86: 5646-5650.
- 25 Ksiezak-Reding H., Liu W.K., Yen S.H. (1992) *Brain Res.* 597: 209-219.
- Lee V., Balin B., Otvos L., Trojanowski J. (1991) A68: a major subunit of paired helical filaments and derivatized forms of normal tau. *Science* 251(4994): 675-8.
- 30 Lewis S., Wang D., Cowan N. (1988) Microtubule-associated protein MAP2 shares a microtubule binding motif with Tau protein. *Science* 242: 936-939.

- Mariani G., Lasku A., Pau A., Villa G., Motta C., Calcagno G., Taddei G.Z., Castellani P., Syrigos K., Dorcaratto A., Epenetos A.A., Zardi L., Viale G.A. (1997) A pilot pharmacokinetic and immunoscintigraphic study with the technetium-99m labelled monoclonal antibody BC-1
5 directed against oncofetal fibronectin in patients with brain tumours. *Cancer* 15: 2484-2489.
- Mohoney F.I., Barthel D.W. (1965) Functional evaluation: The Barthel Index. *Md. State. Med. J.* 24: 61-69.
- 10 Raichle M.E. (1998) Behind the scenes of functional brain imaging: A historical and physiological perspective. *Proc. Natl. Acad. Sci. USA* 95: 765-772.
- Sandrock D., Verheggen R., Helwig A.T., Munz D.L., Markakis E., Emrich D. (1996) Immunoscintigraphy for the detection of brain abscesses. *Nucl. Med. Commun.* 17: 311-316.
- 15 Scandinavian Stroke Study Group (1985) Multicenter trial of hemodilution in ischemic stroke: Background and study protocol. *Stroke* 16: 373-403.
- Selden S., Pollard T. (1983) Phosphorylation of microtubule-associated proteins regulates their
20 interaction with actin filaments. *J. Biol. Chem.* 258(11): 7064-71.
- Tamada K., Fujinaga S., Watanabe R., Yamashita R., Takeuchi Y., Osano M. (1995) Specific deposition of passively transferred monoclonal antibodies against herpes simplex virus type 1 in rat brain infected with the virus. *Microbiol-Immunol.* 39: 861-871
- 25 Vandermeeren M., Mercken M., Vanmechelen E., Six J., Van de Voorde A., Martin J., Cras P. (1993) Detection of tau proteins in normal and Alzheimer's disease fluid with a sensitive sandwich enzyme linked assay *J. Neurochem.* 61: 1828-1834.
- 30 Van de Voorde A., Vanmechelen E., Vandermeeren M., Dessaint F., Beeckman W., Cras P. (1995) Detection of tau in cerebrospinal fluid. *Research Advances in Alzheimer's disease and*

related disorders, Ed. Ibqal, Mortimer, Winblad & Wisniewski, John Wiley & Sons Ltd.

Wakabayashi T., Yoshida J., Okada H., Sugita K., Itoh K., Tadokoro M., Ohshima M. (1995)
Radioimaging of human glioma by indium-111 labelled G-22 anti-glioma monoclonal antibody.

5 Noshuyo-Byori 12: 105-110.

Wilson J.D., Braunwald E., Isselbacher K.J., Petersdorf R.G., Martin J.B., Fauci A.S., Root
R.K. (1991) Harrison's Principles of Internal Medicine, 12th Edition, McGraw-Hill Inc, NY,
USA.

CLAIMS

1. A method for the early detection and/or quantification of CNS damage in an individual, said CNS damage being caused by space-occupying lesions of the CNS, by invasion or metastasis of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents or by a combination of these mechanisms, said method comprising the step of determining the level of tau in said individual and comparing it to the level of tau in control healthy individuals.
2. A method for the early in vitro detection and/or quantification of CNS damage in an individual, said CNS damage being caused by space-occupying lesions of the CNS, by invasion or metastasis of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents or by a combination of these mechanisms, said method comprising the steps of:
 - obtaining a sample from said individual,
 - determining the level of tau in said sample and comparing it to the level of tau in control healthy individuals.
3. A method according to claim 2 in which the sample is taken from the cerebrospinal fluid of the individual.
4. A method according to claim 2 in which the sample is taken from the blood derivatives of the individual.
5. A method according to any of claims 1 to 4 in which the space-occupying lesion of the CNS is a primary brain tumor, benign or malignant, brain metastasis, or a subdural haematoma.
6. A method according to any of claims 1 to 4 in which the invasion or metastasis of the CNS is by leukemia, lymphoma or breast cancer.
7. A method according to any of claims 1 to 4 in which the organisms are bacteria or viruses causing encephalitis or meningitis.

8. A method according to any of claims 1 to 4 in which the anoxia or ischemia is caused by stroke, by cerebral infarction, by cerebral hemorrhage, by thrombosis, by perinatal asphyxia, by Binswanger disease or by vasculitis.

5 9. A method according to any of claims 1 to 4 in which the chemical agent is.

10. A method according to any of claims 1 to 4 in which the physical agent is a trauma, stroke, intracranial pressure or radiation.

10 11. A method according to any of claims 1 to 10 in which CNS damage is detected and/or quantified in order to evaluate the effect of a certain treatment on said CNS damage.

12. The use of tau as an aspecific marker for the manufacture of a diagnostic kit for the early detection and/or quantification in an individual of CNS damage caused by space-occupying lesions of the CNS, by invasion or metastasis of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents, or by a combination of these mechanisms.

15 13. The use of tau as an aspecific marker according to claim 12 in any method according to claims 1 to 11.

20 14. A kit for the early diagnosis of CNS damage in an individual, said CNS damage being caused by space-occupying lesions of the CNS, by invasion of the CNS, by organisms, by anoxia or ischemia, by chemical agents, by physical agents, or by a combination of these mechanisms, comprising a tool for the detection of tau.

25 15. A kit according to claim 14 for use in any method according to claims 1 to 11.

16. A kit according to claims 13 and/or 14 characterised in that said kit comprises:

- a monoclonal antibody (primary antibody) which forms an immunological complex with an epitope of tau;
- a secondary antibody

30

- which can be a monoclonal antibody recognising an epitope of the tau-primary antibody complex, but not recognising the primary antibody alone, or
 - which can be a polyclonal antibody recognising an epitope of the tau-primary antibody complex but not recognising the primary antibody alone, with said polyclonal antibody being preferably purified by immunoaffinity chromatography using immobilized tau or immobilized tau-primary antibody complex;
 - a marker either for specific tagging or coupling with said secondary antibody;
 - appropriate buffer solutions for carrying out the immunological reaction between the primary antibody and the test sample, between the secondary antibody and the tau-primary antibody complex and/or between the secondary antibody and the marker;
 - possibly, for standardisation purposes, a purified protein or a synthetic peptide containing one or more tau epitopes.
17. A method to screen or monitor the effect of compounds which prevent or treat CNS damage comprising the step of determining the level of tau and comparing it to the level of tau in a control sample.

1/3




Figure 1




Figure 2

3/3




Figure 3

WE HEREBY DECLARE THAT ALL STATEMENTS MADE OF OUR OWN KNOWLEDGE ARE TRUE AND THAT ALL STATEMENTS MADE ON INFORMATION AND BELIEF ARE BELIEVED TO BE TRUE; AND FURTHER THAT THESE STATEMENTS WERE MADE WITH THE KNOWLEDGE THAT WILLFUL FALSE STATEMENTS AND THE LIKE SO MADE ARE PUNISHABLE BY FINE OR IMPRISONMENT, OR BOTH, UNDER SECTION 1001 OF TITLE 18 OF THE UNITED STATES CODE AND THAT SUCH WILLFUL FALSE STATEMENTS MAY JEOPARDIZE THE VALIDITY OF THE APPLICATION OR ANY PATENT ISSUED THEREON.

| | |
|---|--|
| Inventor's Full Name <u>1-00 Frank (nmi)</u> <u>HULSTAERT</u> | |
| Inventor's Signature <u>[Signature]</u> | |
| Date: <u>27 FEB 2001</u> | Country of Citizenship: <u>BELGIUM</u> |
| Residence Address | <u>Klaverpoel-15 New: Gontrode Heirweg 44</u> <u>B-9050 Gentbrugge B-9090 MELE</u> <u>BELGIUM BELGIUM</u> <u>BEX</u> |
| Post Office Address, if different from above | <u>same as above</u> |

| | |
|--|---|
| Inventor's Full Name <u>2-00 Eugene (nmi)</u> <u>VANMECHELEN</u> | |
| Inventor's Signature <u>[Signature]</u> | |
| Date: <u>19 Feb 2001</u> | Country of Citizenship: <u>BELGIUM</u> |
| Post Office Address, if different from above | <u>Ten Edestraat 101</u> <u>B-9810 Nazareth Eke</u> <u>BELGIUM</u> <u>BEX</u> <u>same as above</u> |

| | |
|---|--|
| Inventor's Full Name <u>3-00 Hugo (nmi)</u> <u>VANDERSTICHELE</u> | |
| Inventor's Signature <u>[Signature]</u> | |
| Date: <u>15 Feb 2001</u> | Country of Citizenship: <u>BELGIUM</u> |
| Residence Address | <u>Tuinwijklaan 43</u> <u>B-9000 Ghent</u> <u>BELGIUM</u> <u>BEX</u> |
| Post Office Address, if different from above | <u>same as above</u> |

DECLARATION

As below named inventors, we hereby declare that:

Our residence, post office address and citizenship are as stated below next to our names.

The below named inventors are the original, first and joint inventors of the subject matter which is claimed and for which a patent is sought on the invention entitled **TAU AS A MARKER FOR EARLY CNS DAMAGE**, the specification of which was filed as PCT International Application No. **PCT/EP99/06592** on 7 September 1999 and accorded U.S. Serial Number _____.

We hereby state that we have reviewed and understand the contents of the above identified specification, including the claims.

We acknowledge the duty to disclose to the Patent and Trademark Office all information known to us to be material to patentability of the subject matter claimed in this application, as "materiality" is defined in Title 37, Code of Federal Regulations, § 1.56.

We hereby claim foreign priority benefits under Title 35, United States Code, § 119 (a)-(d) of any foreign application(s) for patent listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

PRIOR FOREIGN APPLICATION(S)

Priority Claimed

| | | | |
|-------------------|---------------|--------------------------|------------|
| <u>98870190.0</u> | <u>Europe</u> | <u>08 September 1998</u> | <u>Yes</u> |
| (Number) | (Country) | (Date Filed) | |

We hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code, § 112, we acknowledge the duty to disclose all information known to me to be material to patentability of the subject matter claimed in this application, as "materiality" is defined in Title 37, Code of Federal Regulations, § 1.56, which become available between the filing date of the prior application and the national or PCT international filing date of this application.

| | |
|---------------------------------|-----------------------------|
| <u>PCT/EP99/06592</u> | <u>7 September 1999</u> |
| (International Application No.) | (International Filing Date) |

We hereby direct that all correspondence and telephone calls be addressed to:

Patricia A. Kammerer
Howrey Simon Arnold & White, LLP
750 Bering Drive
Houston, TX 77057-2198
(713) 787-1438

attorneys for the assignee of this application.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:

FRANK HULSTAERT
EUGENE VANMECHELEN
HUGO VANDERSTICHELE

Serial No.:

Filed:

For: **TAU AS A MARKER FOR EARLY CNS
DAMAGE**

§
§
§ Group Art Unit: **Unknown**
§
§ Examiner: **Unknown**
§
§ Atty. Dkt. No.: **11362.0034.PCUS00**
§ **INNS:034---**
§
§ History: **International Application No.:**
§ **PCT/EP99/06592**
§ **Filing Date: 7 September 1999**

ELECTION UNDER 37 C.F.R. §§ 3.71 AND 3.73
AND POWER OF ATTORNEY

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

The undersigned, being Assignee of record of the entire interest in the above-identified application by virtue of an assignment recorded in the United States Patent and Trademark Office as set forth below, hereby elects, under 37 C.F.R. § 3.71, to prosecute the application to the exclusion of the inventors.

The Assignee hereby revokes any previous Powers of Attorney and appoints:

Patricia A. Kammerer, Reg. No. 29,775; M. Todd Rands, Reg. No. 46,249; Matthew L. Madsen, Reg. No. 45,594; M. Suzy Strickland, Reg. No. 45,139; Craig M. Lundell, Reg. No. 30,284; and Janelle D. Waack, Reg. No. 36,300;

each an attorney or agent of the firm of HOWREY SIMON ARNOLD & WHITE, LLP, as its attorney or agent for so long as they remain with such firm, with full power of substitution and revocation, to prosecute the application, to make alterations and amendments therein, to transact all business in the Patent and Trademark Office in connection therewith, and to receive any Letters Patent, and for one year after issuance of such Letters Patent to file any request for a certificate of correction that may be deemed appropriate.

Pursuant to 37 C.F.R. § 3.73, the undersigned has reviewed the evidentiary documents, specifically the Assignment to INNOGENETICS N.V., referenced below, and certifies that to the best of my knowledge and belief, title remains in the name of the Assignee.

Please direct all communications as follows:

Patricia A. Kammerer
HOWREY SIMON ARNOLD & WHITE, LLP
750 Bering Drive
Houston, Texas 77057-2198
(713) 787-1400

ASSIGNEE:
INNOGENETICS N.V.

By: 

Name: Mr. Philippe Archinard
Title: Chief Executive Officer

Date: February 6th, 2001

ASSIGNMENT:

☐ Concurrently filed ☒
☐ Previously recorded

Date:

Reel:

Frames: - -